

Theoretical Physics Institute
University of Minnesota

TPI-MINN-93/32-T
June 1993

Valley Singularities and Baryon Number Violation

Paolo Provero

*Theoretical Physics Institute
University of Minnesota
Minneapolis, MN 55455, USA¹*

Abstract

We consider the valley-method computation of the inclusive cross section of baryon number violating processes in the Standard Model. We show that any physically correct model of the valley action should present a singularity in the saddle point valley parameters as functions of the energy of the process. This singularity prevents the saddle point configuration from collapsing into the perturbative vacuum.

¹e-mail: paolo@physics.spa.umn.edu

Instanton-mediated baryon number violating (BNV) processes in the Standard Model are suppressed, at zero energy, by the factor $e^{-2S_I} \sim e^{-400}$, where S_I is the action of the instanton solution. They were therefore considered as hopelessly unobservable until, in 1990, Ringwald [1] and Espinosa [2] showed that the cross section grows exponentially when energy is raised from zero, because of the increase in available phase space when processes with $O(1/\alpha_W)$ gauge and Higgs bosons in the final state are considered. This observation raised the hope that BNV processes could become observable at energies of order of the sphaleron mass ~ 10 TeV. Despite a lot of theoretical effort (for a review see Ref. [3]), a reliable computation of the cross section at the relevant energies is still lacking.

It is customary to express the leading term in semiclassical approximation to the BNV cross section as

$$\sigma_{BNV}(E) \sim \exp(2S_I F_{hg}(E)) \quad (1)$$

where E is the energy of the process and $F_{hg}(E)$ is the so-called holy-grail function.

It is by now well known how to express $F_{hg}(E)$ as a series expansion in $(E/E_0)^{2/3}$, where E_0 is an energy scale of the order of the sphaleron mass. Choosing

$$E_0 = \frac{\sqrt{6}\pi M_W}{\alpha_W} \sim 17 \text{ TeV} \quad (2)$$

the known terms in the expansion of F_{hg} are [4]

$$\begin{aligned} F_{hg}(E) = & -1 + \frac{9}{8} \left(\frac{E}{E_0}\right)^{4/3} - \frac{9}{16} \left(\frac{E}{E_0}\right)^{6/3} \\ & + \frac{3}{32} \left(4 - 3\frac{M_H^2}{M_W^2}\right) \left(\frac{E}{E_0}\right)^{8/3} \log\left(\frac{E}{E_0}\right) + O\left(\frac{E}{E_0}\right)^{8/3} \end{aligned} \quad (3)$$

What is needed is a reliable way of extending the computation to finite values of the ratio E/E_0 , and therefore predict if BNV processes can become observable at energies accessible to the next generation of colliders.

The valley method [5, 6] provides in principle such an analytic continuation of the expansion (3), in the (somewhat ill-defined) approximation where initial-state corrections are neglected. σ_{BNV} is computed, using the optical theorem, as the imaginary part of the forward elastic scattering (FES) amplitude of two fermions going into two fermions through the background

field of a deformed instanton–antiinstanton ($I\bar{I}$) pair. The cross section is expressed, to leading order in semiclassical approximation, as

$$\sigma_{BNV}(E) \sim \text{Im} \int dR d\rho_I d\rho_{\bar{I}} \exp(ER - S_v(R, \rho_I, \rho_{\bar{I}})) \quad (4)$$

where $S_v(R, \rho_I, \rho_{\bar{I}})$ is the classical action along the valley trajectory, which connects a far separated $I\bar{I}$ pair with the perturbative vacuum. The valley trajectory is chosen in such a way that the functional integration along the directions orthogonal to it is gaussian, and can therefore be neglected in the leading order of semiclassical approximation. The trajectory is parametrized by the distance R between the centers of the (deformed) I and $\bar{I}$ and their radii $\rho_I, \rho_{\bar{I}}$. The integral is computed in saddle-point approximation.

Unfortunately, the valley action in the Standard Model is not known beyond an expansion in powers of ρ/R and ρ (from now on we anticipate that at the saddle point $\rho_I = \rho_{\bar{I}} \equiv \rho$). The expression of F_{hg} derived from this valley action is the expansion (3). In general, it is possible to prove [7] that the valley method computation of $F_{hg}(E)$ as an expansion in E/E_0 is equivalent to all orders to the direct computation in the one-instanton sector. This suggests that $F_{hg}(E)$ can be correctly computed with the valley method for all energies in which $F_{hg}(E)$ is an analytic function of E .

An approximate model for the electroweak valley action has been proposed in [6], where the (exactly known) QCD valley [8] is supplemented with a conformal symmetry breaking term

$$S_{csb} = 2\pi^2 v^2 \rho^2 \quad (5)$$

where v is the VEV of the Higgs field. This means that the $I\bar{I}$ interaction in the Higgs sector is completely neglected, the rationale being that Higgs boson production in the final state is subdominant with respect to gauge boson production at $E \ll E_0$. In this model, the holy-grail function hits the unitarity limit $F_{hg} = 0$ at a finite energy $E_{KR} \sim 45\text{TeV}$, and the cross section loses its exponential suppression. The saddle point configuration collapses into the perturbative vacuum as $E \rightarrow E_{KR}$, and $F_{hg}(E)$ does not show any singularity² between $E = 0$ and $E = E_{KR}$.

In this Letter we will show that the absence of singularities and the collapse into the perturbative vacuum are artifacts of the approximation defining the model, and are bound to disappear in a qualitatively realistic

²As already pointed out in Ref.[9], the singularity found in Ref. [10] in the Khoze–Ringwald model is an artifact of a particular way of solving the saddle point equations.

model of the Higgs contribution to the valley action. In fact we will show that the existence of a singularity preventing the collapse to the perturbative vacuum is an unavoidable feature of any realistic model of the valley action, which stems from general features of the valley trajectory.

The saddle point evaluation of the integral (4) gives for the holy-grail function the expression

$$F_{hg}(E) = ER_*(E) - S_v(R_*(E), \rho_*(E)) \quad (6)$$

where $R_*(E)$, $\rho_*(E)$ are solution of the saddle point equations

$$E = \frac{\partial S_v}{\partial R} \quad (7)$$

$$0 = \frac{\partial S_v}{\partial \rho} \quad (8)$$

From Eq. (7) we can immediately see that the perturbative vacuum cannot be the saddle point for any nonzero energy. Indeed, the action must be stationary for all deformations of the fields around the perturbative vacuum, and in particular it must be

$$\left. \frac{\partial S_v}{\partial R} \right|_{\text{pert.vacuum}} = 0 \quad (9)$$

Note that the perturbative vacuum cannot even be the limit of the saddle point configurations for $E \rightarrow E_{lim} \neq 0$, because it is impossible to have values of the action arbitrarily close to zero while keeping $\partial S/\partial R$ finite and nonzero.

It is interesting to see how this seemingly impossible phenomenon occurs in the Khoze–Ringwald model of Ref. [6]. In this model

$$\frac{\partial S_v}{\partial R} = \int d^4x \frac{\delta S_v}{\delta A_\mu^a} \frac{\partial A_\mu^a}{\partial R} \quad (10)$$

and

$$\lim_{E \rightarrow E_{KR}} \frac{\delta S_v}{\delta A_\mu^a} = 0 \quad (11)$$

However, the field configuration [8] is given by ³

$$A_\mu^a(x) = \frac{2}{g} \eta_{a\mu\nu} x_\nu \left\{ \frac{1}{x^2 + \rho^2/z} - \frac{1}{x^2 + \rho^2 z} \right\} \quad (12)$$

³We consider here the concentric $I\bar{I}$ configuration, from which the non-concentric one can be obtained by a coordinate inversion, see Ref. [8].

where

$$z = 1 + \frac{\xi^2}{2} + \xi \sqrt{1 + \frac{\xi^2}{4}}, \quad \xi = \frac{R}{\rho} \quad (13)$$

so that

$$\frac{\partial A_\mu^a}{\partial R} = \frac{2}{g} \eta_{a\mu\nu} \rho x_\nu \left\{ \frac{1}{(x^2 z + \rho^2)^2} + \frac{1}{(x^2 + \rho^2 z)^2} \right\} \frac{dz}{d\xi} \quad (14)$$

Being

$$\lim_{E \rightarrow E_{KR}} \rho_*(E) = 0 \quad (15)$$

$$\lim_{E \rightarrow E_{KR}} z_*(E) = 1 \quad (16)$$

$\partial A_\mu^a / \partial R$ develops a singularity in $x = 0$ for $E \rightarrow E_{KR}$, which is responsible for the non-vanishing of the integral (10).

Returning to the general case, we note that the saddle point equations (7) and (8) can be solved to give $R_*(E)$ and $\rho_*(E)$ whenever

$$W = \det \begin{pmatrix} \frac{\partial^2 S_v}{\partial R^2} & \frac{\partial^2 S_v}{\partial R \partial \rho} \\ \frac{\partial^2 S_v}{\partial \rho \partial R} & \frac{\partial^2 S_v}{\partial \rho^2} \end{pmatrix} \neq 0 \quad (17)$$

Now for a far-separated $I\bar{I}$ we have $W < 0$, because this is an unstable (asymptotical) solution of the equations of motion⁴. This is precisely the reason why a far separated $I\bar{I}$ pair contributes to the *imaginary* part of the FES amplitude. On the other hand, in the perturbative vacuum, which is a stable solution of the equations of motion, we have $W > 0$. It follows the existence of a boundary in (R, ρ) space separating the $W > 0$ and $W < 0$ regions. On this boundary $W = 0$.

Now consider the line defined in (R, ρ) space by the solutions of Eq.(8). It joins the far-separated $I\bar{I}$ pair with the perturbative vacuum, which are both solutions of (8); it follows that this line will intersect the $W = 0$ boundary in a point (R_i, ρ_i) . Here Eqs. (7) and (8) cannot be solved, and the saddle point parameters $R_*(E)$, $\rho_*(E)$ develop a singularity in

$$E_s = \frac{\partial S_v}{\partial R}(R_i, \rho_i) \quad (18)$$

⁴More precisely, $W \rightarrow 0^-$ when $R \rightarrow \infty$.

The singularity of the field derivative in the Khoze–Ringwald model is again responsible for the absence of such a singularity, because our mechanism is based on the fact that the vacuum is a stationary point of the action. On the other hand, it is not surprising that in the $O(3)$ σ -model studied in Ref. [9], where the conformal symmetry breaking term in the action is derived from a true Lagrangian, a singularity is found in the saddle point parameters: in this point the saddle point equations (7) and (8) cannot be inverted to give $R_*(E)$ and $\rho_*(E)$ and this *implies* that the $W = 0$ boundary has been reached (if $W \neq 0$ the saddle point equations can certainly be solved).

In summary, we have described a general mechanism giving rise to singularities of the saddle point valley parameters for any model of the valley action displaying the correct qualitative behaviour, in the Standard Model as well as in all the toy models developed to gain insight in the problem (our argument is independent on the number of parameters needed to describe the valley trajectory, and is actually even more immediate for one-parameter models like the two-dimensional Abelian Higgs Model). This mechanism necessarily prevents the saddle point configuration from collapsing into the perturbative vacuum. In conclusion, let us remark the following:

- (i) Our explanation of the origin of the singularities is different from the one proposed in Ref. [11], because it is independent from the choice of the weight function defining the valley trajectory, and relies only on the general features of the valley trajectory.
- (ii) Our scenario is too general to give indications about the value of the holy-grail function at the singularity point: in principle the singularity can occur after the cross section has lost its exponential suppression. However, in the $O(3)$ σ -model of Ref. [9], the singularity occurs, when the parameters of the model are chosen in a self-consistent way, at an energy where the cross-section is still exponentially suppressed.
- (iii) A related problem is the relevance of the FES scattering amplitude computed in the valley background to baryon number violation [12, 13, 14, 15, 16]: roughly speaking the problem is that when the I and $\bar{I}$ start to overlap the intermediate states of the FES amplitude (mapped by unitarity in the final states of the BNV process) do not have anymore the baryon number predicted by the anomaly law; the danger is to mistakenly consider baryon number *conserving* contributions to the cross section. Let us notice

that in our scenario the problem becomes less severe, because the complete overlapping of I and $\bar{I}$ is prevented by the singularity.

Acknowledgements

The author would like to thank M. Shifman for many useful conversations. Discussions with R. Guida, K. Konishi, M. Maggiore and N. Magnoli are also gratefully acknowledged. This work is supported by an INFN grant.

References

- [1] A. Ringwald, *Nucl. Phys.* **B330**,1 (1990).
- [2] O. Espinosa, *Nucl. Phys.* **B343**, 310 (1990).
- [3] M. Mattis, *Phys. Rep.* **214**, 159 (1992).
- [4] I. Balitsky and A. Schäfer, preprint UFTP 323/1992 and PSU/TH/119, hep-ph 9304261.
- [5] V.V. Khoze and A. Ringwald, *Nucl. Phys.* **B355**, 351 (1991).
- [6] V.V. Khoze and A. Ringwald, *Phys. Lett.* **B259**, 106 (1991).
- [7] P. Arnold and M. Mattis, *Phys. Rev.* **D44**, 3650 (1991).
- [8] A. Yung, *Nucl. Phys.* **B297**, 47 (1988).
- [9] J. Grandy and M. Mattis, preprint LA-UR-93-811, hep-ph-9303218.
- [10] E. Shuryak and J. Verbaarschot, *Phys. Rev. Lett.* **68**, 2576 (1992).
- [11] N. Dorey and M. Mattis, *Phys. Lett.* **B277**, 337 (1992).
- [12] M. Maggiore and M. Shifman, *Phys. Rev.* **D47**, 3610 (1993).
- [13] I. Kogan, M. Shifman and A. Vainshtein, unpublished.
- [14] R. Guida and K. Konishi, *Phys. Lett.* **B298**, 371 (1993).
- [15] R. Guida and K. Konishi, preprint GEF-Th-8/1993, hep-ph 9303232.
- [16] R. Guida, K. Konishi and N. Magnoli, preprint GEF-Th-14/1993, hep-ph 9306284.